



北海道大学
HOKKAIDO UNIVERSITY



広島大学



熊本大学
Kumamoto University



The Genotype to Phenotype Japan (G2P-Japan)

WPI-ICReDD

B.1.1.529, BA

2 5

The Genotype to Phenotype Japan (G2P-Japan)

VOC variant of concern

B.1.1.529, BA

2 5

Nature

2022 2 1

500 SARS-CoV-2 2022 1 3
2019
2020
B.1.1.529, BA 11 26
2022 1

2 5

5

G2P-Japan

G2P-Japan

genotype
phenotype

AMED
20fk0108413 20fk0108451
CREST JPMJCR20H4

Nature 2 1

Attenuated fusogenicity and pathogenicity of SARS-CoV-2 Omicron variant

#, #, #, #, #, #, , Hesham Nasser,
, , , , , , , , ,
, , , The Genotype to Phenotype Japan (G2P-Japan) Consortium, ,
, , *, *, *, *

#Equal contribution; *Corresponding author

DOI: 10.1038/s41586-022-04462-1

URL: <https://www.nature.com/articles/s41586-022-04462-1>

03-6409-2212

FAX 03-6409-2213

keisato@g.ecc.u-tokyo.ac.jp

090-9832-9760

koho@ims.u-tokyo.ac.jp

011-706-2610 FAX 011-706-2092

jp-press@general.hokudai.ac.jp

096-342-3269

sos-koho@jimu.kumamoto-u.ac.jp

082-424-3701 FAX 082-424-6040

koho@office.hiroshima-u.ac.jp


VOC variant of concern

VOC variant of concern

B.1.1.7
B.1.617.2, AY

B.1.351
B.1.1.529, BA


P.1



VeroE6/TMPRSS2

48

N
B.1.1



B.1.1

Penh, Rpef,

[SpO2]

P

